



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### **SUMOylation of phytochrome-B negatively regulates light-induced signaling in *Arabidopsis thaliana***

**Citation for published version:**

Sadanandom, A, Ádám, É, Orosa, B, Viczián, A, Klose, C, Zhang, C, Josse, EM, Kozma-Bognár, L & Nagy, F 2015, 'SUMOylation of phytochrome-B negatively regulates light-induced signaling in *Arabidopsis thaliana*', *Proceedings of the National Academy of Sciences (PNAS)*, vol. 112, no. 35, pp. 11108-11113.  
<https://doi.org/10.1073/pnas.1415260112>

**Digital Object Identifier (DOI):**

[10.1073/pnas.1415260112](https://doi.org/10.1073/pnas.1415260112)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Publisher's PDF, also known as Version of record

**Published In:**

Proceedings of the National Academy of Sciences (PNAS)

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# SUMOylation of phytochrome-B negatively regulates light-induced signaling in *Arabidopsis thaliana*

Ari Sadanandom<sup>a,1</sup>, Éva Ádám<sup>b</sup>, Beatriz Orosa<sup>a</sup>, András Viczián<sup>b</sup>, Cornelia Klose<sup>c</sup>, Cunjin Zhang<sup>a</sup>, Eve-Marie Josse<sup>d</sup>, László Kozma-Bognár<sup>b</sup>, and Ferenc Nagy<sup>b,d,1</sup>

<sup>a</sup>School of Biological and Biomedical Sciences, University of Durham, Durham DH1 3LE, United Kingdom; <sup>b</sup>Plant Biology Institute, Biological Research Centre, H-6726 Szeged, Hungary; <sup>c</sup>Institute of Botany, University of Freiburg, D-79104 Freiburg, Germany; and <sup>d</sup>Institute of Molecular Plant Science, School of Biology, University of Edinburgh, Edinburgh EH9 3JR, United Kingdom

Edited by George Coupland, Max Planck Institute for Plant Breeding Research, Cologne, Germany, and approved July 16, 2015 (received for review August 8, 2014)

The red/far red light absorbing photoreceptor phytochrome-B (phyB) cycles between the biologically inactive (Pr,  $\lambda_{\max}$ , 660 nm) and active (Pfr;  $\lambda_{\max}$ , 730 nm) forms and functions as a light quality and quantity controlled switch to regulate photomorphogenesis in *Arabidopsis*. At the molecular level, phyB interacts in a conformation-dependent fashion with a battery of downstream regulatory proteins, including PHYTOCHROME INTERACTING FACTOR transcription factors, and by modulating their activity/abundance, it alters expression patterns of genes underlying photomorphogenesis. Here we report that the small ubiquitin-like modifier (SUMO) is conjugated (SUMOylation) to the C terminus of phyB; the accumulation of SUMOylated phyB is enhanced by red light and displays a diurnal pattern in plants grown under light/dark cycles. Our data demonstrate that (i) transgenic plants expressing the mutant phyB<sup>Lys996Arg</sup>-YFP photoreceptor are hypersensitive to red light, (ii) light-induced SUMOylation of the mutant phyB is drastically decreased compared with phyB-YFP, and (iii) SUMOylation of phyB inhibits binding of PHYTOCHROME INTERACTING FACTOR 5 to phyB Pfr. In addition, we show that OVERLY TOLERANT TO SALT 1 (OTS1) de-SUMOylates phyB in vitro, it interacts with phyB in vivo, and the *ots1/ots2* mutant is hyposensitive to red light. Taken together, we conclude that SUMOylation of phyB negatively regulates light signaling and it is mediated, at least partly, by the action of OTS SUMO proteases.

photoreceptor | phytochrome | sumoylation | signaling | photomorphogenesis

Plants are sessile organisms; thus, they have to adapt to the ever-changing environment by modifying their growth and developmental programs. To respond to changes in ambient light conditions, plants evolved a battery of photoreceptors including the blue/UVA absorbing cryptochromes and phototropins (1, 2), the red/far red absorbing phytochromes (phys) (3), and the UVB-specific photoreceptor UVR8 (4). The red light/far red light (RL/FRL) absorbing phys exist as dimers, and each monomer has a covalently linked open tetrapyrrole chain as chromophore. They are synthesized in their biologically inactive Pr form (RL-absorbing state;  $\lambda_{\max}$ , 660 nm) and converted into the biologically active Pfr (FRL-absorbing state;  $\lambda_{\max}$ , 730 nm) by RL. Subsequent FRL treatment converts the Pfr form back into Pr. The Pr and Pfr conformers have partially overlapping absorption spectra; thus, phys cycle between their Pfr/Pr forms, and the ratio of Pr/Pfr forms is determined by the RL/FRL content of the incipient sunlight (5). In *Arabidopsis* five phys have been identified (phyA, phyB, phyC, phyD, and phyE), and among these, phyB has been shown to be especially important after seedling establishment (6).

The overwhelming majority of molecular events underlying phyB-controlled photomorphogenesis take place in the nucleus. Light in a quality- and quantity-dependent fashion induces translocation of phyB Pfr in the nuclei (7, 8), where it interacts with downstream acting regulatory proteins including PHYTOCHROME-INTERACTING-FACTORS (PIFs) (9). The very early

steps of phyB signaling include (i) inactivation or alteration of the substrate specificity of CONSTITUTIVE PHOTOMORPHOGENIC 1 (COP1) that targets proteins to degradation (10), (ii) degradation and/or modulation of the transcriptional activity of negative regulatory PIF TFs (11), and (iii) induction of transcriptional cascades that modulate the expression of 2,500–3,000 genes of the *Arabidopsis* genome (12).

The number of phyB Pfr molecules quantitatively determines physiological responses. Thus, it is evident that any molecular process that modifies (i) the abundance of phyB Pfr such as ubiquitination of the photoreceptor followed by its subsequent degradation (13, 14), (ii) the accessibility (8), and (iii) the capacity of phyB Pfr to interact with its signaling partners alters the flux of RL-induced signaling. Very recently it was shown that the Ser86 (15) and/or Tyr104 amino acid residues (16) of phyB are phosphorylated *in planta*. These reports established that phosphorylation of S86 accelerates thermal relaxation (light-independent reversion of Pfr into Pr, also called dark reversion) of phyB, whereas light-induced phosphorylation of Tyr104 inhibits binding of PIF3 to phyB Pfr. Taken together, these results clearly demonstrated that multiple phosphorylation of phyB negatively regulates the action of this photoreceptor.

Reversible conjugation of small ubiquitin-like modifier (SUMO) to the target proteins represents an emerging posttranslational modification (17–20). Work performed in mammalian and yeast cells showed that (i) reversible SUMOylation of proteins regulates a

## Significance

The photoreceptor phytochrome-B (phyB) cycles between its active Pfr [far red light (FRL)-absorbing state  $\lambda_{\max}$ , 730 nm] and inactive Pr [red light (RL)-absorbing state  $\lambda_{\max}$ , 660 nm] forms and regulates as red/far red light-activated/inactivated molecular switch plant growth and development. Here we show that conjugation of small ubiquitin-like modifier to the photoreceptor inhibits interaction of phyB Pfr with its immediate signaling partner PHYTOCHROME INTERACTING FACTOR 5 (PIF5). The impaired interaction of these proteins negatively affects photomorphogenic responses; thus, SUMOylation similar to phosphorylation plays a role in desensitizing phyB-mediated signaling. OVERLY TOLERANT TO SALT 1 and 2 (OTS1 and OTS2) are involved in regulating phyB action as these SUMO proteases mediate deconjugation of SUMO from phyB.

Author contributions: A.S. and F.N. designed research; A.S., É.A., B.O., A.V., C.K., C.Z., E.-M.J., and L.K.-B. performed research; A.S. and F.N. analyzed data; and A.S. and F.N. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

<sup>1</sup>To whom correspondence may be addressed. Email: nagyf@brc.hu or ari.sadanandom@durham.ac.uk.


This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1415260112/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1415260112/-DCSupplemental).

broad spectrum of cellular processes and (ii) addition of SUMO either modifies the activity of the target protein and/or promotes interaction with specific proteins harboring the SUMO-interacting motif (21, 22). It has also been established that in plants, like in mammalian and yeast cells, a SUMOylation pathway exists for SUMOylation and de-SUMOylation of proteins (23, 24). More specifically, it was shown that in *Arabidopsis thaliana* (i) expression of isoforms of SUMO1, -2, -3, and -5 can be detected (25); (ii) conjugation of SUMO to target proteins occurs via the same sequential biochemical steps as in yeast (26, 27); (iii) these reactions are catalyzed by the E1 heterodimer (SAE1a/b and SAE2), a single E2 (SCE1), and two E3 (SIZ1 and MMS2/HPY2) enzymes as well as E4-type SUMO ligases (28); whereas (iv) de-conjugation of SUMO substrates is performed by ubiquitin-like SUMO-specific proteases including ESD4 and OTS1/2 (29–34). It appears that in plants reversible conjugation of SUMO to target proteins is particularly important in regulating responses to abiotic environmental stresses including heat shock (35), phosphate starvation (36), drought (37), and salt stress (31), but SUMOylation of specific proteins also modifies responses to ABA signaling (38) and phytopathogen infection (39) and modulates flowering time (33).

Light is arguably the most important environmental factor regulating plant growth and development, yet until now, in vitro and in planta studies have not indicated SUMOylation of any component of photoreceptor-controlled light signaling cascades. Here we report that (i) the photoreceptor phyB is reversibly SUMOylated in planta, (ii) SUMOylated phyB preferentially accumulates in light, (iii) binding of PIF5 to the SUMOylated phyB Pfr is significantly reduced, and (iv) transgenic plants expressing the mutant phyB<sup>Lys996Arg</sup> photoreceptor are hypersensitive to RL. Furthermore, we show that the *ots1/ots2* mutant displays hypersensitivity to RL and enhanced phyB SUMOylation, whereas OTS1 de-SUMOylates phyB in vitro and binds to phyB in planta. Taken together, these data suggest that OTS1/OTS2 is involved in mediating the reversible SUMOylation of phyB and that SUMOylation of phyB desensitizes RL-induced signaling, similarly to phosphorylation of the photoreceptor.

## Results

**PhyB Is SUMOylated in Plants Grown in Light/Dark Cycles.** Experimentally determined SUMO attachment sites contain the ΨKXE sequence, where Ψ (Psi) is a large hydrophobic amino acid and K represents the lysine that binds SUMO (40). The ever-increasing number of identified SUMOylation sites in eukaryotic cells made it possible to construct reliable sequence analysis programs to identify, in silico, sites that are potential targets for SUMOylation in vivo (41). We used the SUMOsp 2.0 software to search for potential SUMOylation site(s) in phyB of *A. thaliana* (42). This search predicted the presence of two potential SUMOylation sites, K996 (high probability) in the C terminus and Lys475 (low probability) in the N terminus of the photoreceptor in AtphyB (Table S1). Fig. 1A shows the sequence context for the Lys996 in *Arabidopsis* as well as in phyB sequences of other plants species. To test if phyB is indeed SUMOylated in planta, first we analyzed SUMOylation of AtphyB-YFP in *Nicotiana benthamiana* leaves transiently coexpressing HA-SUMO and phyB-YFP or the phyB<sup>Lys475Arg</sup> or phyB<sup>Lys996Arg</sup> mutants. Our data demonstrate that AtphyB is SUMOylated and that mutation of Lys996 but not that of Lys475 inhibits SUMOylation of the photoreceptor in tobacco leaves (Fig. S1). To confirm that phyB is indeed subject of SUMOylation, we grew transgenic *phyB-9 Arabidopsis* plants expressing the 35S:PHYB-YFP transgene in light/dark (L/D) cycles, isolated the phyB-YFP fusion protein by affinity purification, and determined its SUMOylation status by using an antibody specific to AtSUMO-1. Accumulation of SUMOylated phyB-YFP reached high levels in 12 h RL/12 h D (Fig. 1B), and it was around the limit of detection in 12 h FRL/12 h D cycles (Fig. 1C). This figure also demonstrates that




**Fig. 1.** The photoreceptor phyB is SUMOylated in seedlings grown in RL/D cycles. (A) Analysis of *A. thaliana* (At, NP\_179469), *Solanum lycopersicum* (Sl, XP\_010318997), *Oryza sativa* (Os, EEE58928), *Triticum aestivum* (Ta, BAP91202), and *Zea mays* (Zm, NP\_001168077) phyB protein sequences by SUMOsp 2.0 software identified sequence motifs as potential targets for SUMOylation. The Lys996 amino acid of the conserved SUMOylation site for AtphyB is indicated by an arrow. Numbers at the beginning and at the end of the presented sequences indicate the corresponding amino acid positions, and numbers in brackets show the length of PHYB proteins from different species. (B) *phyB-9* seedlings expressing the 35S:PHYB-GFP transgene were grown on MS medium in 12 h RL (25  $\mu\text{mol m}^{-2}\text{s}^{-1}$ )/12 h D cycles and 12 h FRL (10  $\mu\text{mol m}^{-2}\text{s}^{-1}$ )/12 h D cycles for 5 d. Samples were harvested on day 6, at MOD and MON. phyB-GFP was immunoprecipitated by using GFP-Trap agarose beads (IP:αGFP), and samples containing identical amounts of the fusion protein were analyzed. phyB-GFP was detected by anti-GFP (IB:αGFP), whereas AtSUMO1-conjugated phyB-GFP was visualized using anti-SUMO1 antibody (IB:αSUMO1).

accumulation of the SUMOylated phyB in RL/D cycles is light regulated, as it accumulates to higher levels at the middle of day (MOD) and decreases to lower levels by the middle of the night (MON). The abundance of phyB does not significantly change in the 35S:PHYB-YFP plants grown under these conditions (Fig. S2A); thus, we conclude that phyB is SUMOylated in a reversible fashion in RL/D cycles. We also monitored SUMOylation of fusion proteins containing truncated forms of phyB including phyB(1–651)-YFP (43) and phyB(1–991)-YFP. The abundance of the truncated fusion proteins is similar to that of phyB-GFP, yet SUMOylation of these fusion proteins was below detection level in plants grown in RL/D cycles (Fig. S2B). Taken together, these data show that phyB is SUMOylated in vivo, the SUMOylated form of phyB accumulates to high levels in RL, and in harmony with data obtained by the transient SUMOylation assays, the site of reversible SUMOylation of phyB is located at the C-terminal domain of the photoreceptor.

**Transgenic phyB-9 plants expressing the mutant phyB<sup>Lys996Arg</sup>-YFP are hypersensitive to RL.** To determine the physiological importance of the predicted SUMOylation site in phyB, we generated transgenic lines expressing, under the control of the viral 35S promoter, the phyB-GFP and phyB<sup>Lys996Arg</sup>-YFP fusion proteins at comparable levels (Fig. S2C, Set 1). Analysis of photomorphogenic responses in RL demonstrated that the transgenic 35S:PHYB-GFP lines displayed characteristic overexpression phenotypes compared with wild-type (Col-0) seedlings including fluence rate-dependent inhibition of hypocotyl elongation (Fig. 2A and Fig. S3A). Importantly, we also show that the 35S:PHYB<sup>Lys996Arg</sup>-YFP lines displayed greater hypersensitivity to continuous RL not only for hypocotyl elongation (Fig. 2A) but cotyledon expansion (Fig. 2C) and cotyledon opening (Fig. 2D), indicating enhanced signaling by the mutant photoreceptor. In contrast, these lines did not display exaggerated responsiveness to FRL or blue light (BL) (Fig. S3C and D). To recapitulate






**Fig. 2.** Seedlings expressing the mutant phyB<sup>Lys996Arg</sup>-YFP are hypersensitive to RL. Relative fluence rate-dependent inhibition of hypocotyl elongation of wild-type Col-0 (black square), phyB-9 (star symbol), and transgenic phyB-9 plants expressing phyB-GFP (gray square) or phyB<sup>Lys996Arg</sup>-YFP (white diamond) under the control of the constitutively active (A) viral 35S promoter or (B) the *Arabidopsis* LIP1 promoter is shown. Seedlings were grown in darkness or at the indicated fluence rates of RL for 4 d. Hypocotyl lengths of irradiated seedlings were measured, and the average value of 25–30 seedlings was calculated and normalized to the hypocotyl length of D-grown plants. (C) Transgenic phyB-9 seedlings expressing 35S:PHYB-GFP (black columns) or 35S:PHYB<sup>Lys996Arg</sup>-YFP (striped columns) were grown at the indicated fluence rates of RL for 4 d. Cotyledon areas of irradiated seedlings were measured, and the average value was normalized to the cotyledon area of D-grown plants. (D) Cotyledon angle of D- or RL-grown seedlings analyzed in Fig. 2C was measured. All experiments shown in Fig. 2 were repeated three times. Error bars indicate SE. Asterisks show statistically significant differences between phyB-GFP or phyB<sup>Lys996Arg</sup>-YFP expressor plants as determined by Student's *t* test (\**P* < 0.05; \*\**P* < 0.01; \*\*\**P* < 0.001).

these data in plants containing phyB at physiological levels, we generated transgenic phyB-9 plants that expressed phyB-YFP or phyB<sup>Lys996Arg</sup>-YFP fusion proteins under the control of the moderately active LIP1 (*LIGHT INSENSITIVE PERIOD 1*) promoter (44). Expression of the LIP1:PHYB-YFP transgene did not result in an overexpression phenotype but restored RL fluence rate-dependent inhibition of hypocotyl elongation similar to the wild-type (Col-0) level (Fig. 2B). Furthermore, we found that LIP1:PHYB<sup>Lys996Arg</sup>-YFP seedlings are also hypersensitive to RL (Fig. 2B and Fig. S3B) compared with LIP1:PHYB-YFP, although the abundance of the fusion proteins is identical (Fig. S2C). Thus, we conclude that the mutant phyB<sup>Lys996Arg</sup>-YFP, overexpressed or expressed at the physiological level, invariably displays enhanced RL-induced signaling. To test if enhanced signaling by the mutant phyB<sup>Lys996Arg</sup>-YFP was brought about by increased/accelerated translocation of the mutant photoreceptor into the nucleus, we determined RL-dependent nuclear accumulation of the various phyB fusion proteins *in planta*. The data obtained indicate that the velocity of import into the nuclei and the formation as well as the size of phyB-YFP and phyB<sup>Lys996Arg</sup>-YFP-associated photobodies do not differ *in planta* (Fig. S4).


**The Lys<sup>996</sup>Arg mutation reduces accumulation of SUMOylated phyB-YFP.** To test if enhanced RL-induced signaling of the mutant phyB<sup>Lys996Arg</sup>-YFP photoreceptor was indeed brought about by its altered SUMOylation, we performed two lines of experiments.

First we determined whether the Lys/Arg substitution affected rates of photoconversion and/or dark reversion of phyB. Our data demonstrate that the basic photochemical properties of the mutant phyB do not significantly differ from those of phyB (Fig. S5). Next we compared the accumulation of SUMOylated phyB-GFP and phyB<sup>Lys996Arg</sup>-YFP *in planta*. We found that amounts of detectable SUMOylated phyB-GFP and phyB<sup>Lys996Arg</sup>-YFP are low in etiolated seedlings, whereas RL treatment elevates the level of SUMOylated phyB-GFP but not that of phyB<sup>Lys996Arg</sup>-YFP (Fig. 3). Accumulation of SUMOylated phyB-GFP increases slowly during the first 3–6 h of irradiation and reaches a substantially higher level after 24 h of treatment with RL (Fig. 3), and elevated intensities of RL are more effective to induce accumulation of SUMOylated phyB-GFP (Fig. S6A). Exposure of etiolated seedlings to white light (WL) also differentially modulates accumulation of SUMOylated phyB-GFP and phyB<sup>Lys996Arg</sup>-YFP (Fig. 4). Importantly, this figure also illustrates that the constitutively active phyB<sup>Tyr276His</sup>-YFP photoreceptor is hyper-SUMOylated in a light-independent fashion in etiolated seedlings. In addition, we found that nuclear pool of phyB is more prone to SUMOylation by comparing the level of SUMOylation of phyB in phyB-GFP-, phyB-GFP-nuclear localization signal (NLS)-, and phyB-GFP-nuclear export signal (NES)-expressing transgenic lines (Fig. S6B). Taken together, these data suggest that SUMOylated phyB preferably accumulates in its Pfr form in the nucleus, its accumulation is RL fluence rate-dependent, and perturbation of the putative SUMOylation site significantly reduces the detectable amount of SUMOylated phyB-GFP Pfr leading to enhanced signaling.


**Binding of PIF5 to phyB Pfr is inhibited by SUMOylation of the photoreceptor.** PIF1 and PIF3–PIF8 represent a subgroup of the bHLH (basic-helix–loop–helix) type transcription factor family that binds to phyB Pfr (45). PIFs negatively regulate signaling by phyB (46, 47) and function as a signaling hub to integrate light and hormone-induced signaling cascades (9, 48). Binding of PIFs to phyB initiates phosphorylation and subsequent degradation (49) or inactivation of these TFs (11, 50) and results in altered transcription of hundreds of genes in seedlings exposed to RL. It has been shown that compromised binding of PIFs to phyB Pfr results in reduced sensitivity to RL (49). To test whether SUMOylation of phyB interferes with PIF5 binding to phyB Pfr, we performed pull-down assays from infiltrated *N. benthamiana* leaves transiently coexpressing phyB-YFP or phyB<sup>Lys996Arg</sup>-YFP with PIF5-MYC and HA-SUMO1 proteins. Our data demonstrate that binding of PIF5 to phyB-GFP (Fig. 5, lane 1) is reduced compared with phyB<sup>Lys996Arg</sup>-YFP (Fig. 5, lane 2), although the amount of phyB-YFP and phyB<sup>Lys996Arg</sup>-YFP is identical. This figure also demonstrates that Lys996Arg substitution inhibits SUMOylation of the receptor. We also note that in these assays the level of SUMOylated phyB-YFP is likely to be lowered by conjugation of *N. benthamiana* SUMO1. Taken together, these results suggest that binding of PIF5 to phyB Pfr is inhibited by the SUMOylation of the photoreceptor.



**Fig. 3.** RL-induced accumulation of the SUMOylated phyB<sup>Lys996Arg</sup> mutant photoreceptor is reduced. phyB-9 seedlings expressing phyB-GFP (WT) or phyB<sup>Lys996Arg</sup>-YFP (Lys996Arg) fusion proteins were grown in darkness for 4 d (D) and subsequently transferred to RL ( $12.5 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) for the indicated time. Sample preparation and detection of SUMOylation were performed as described in Fig. 1.







**Fig. 6.** The SUMO proteases OTS1 and OTS2 mediate de-SUMOylation of phyB. (A) The *ots1/ots2* mutant is hypersensitive to RL. Fluence rate-dependent relative inhibition of hypocotyl elongation of Col-0 (black square), *phyB-9* (star symbol), *ots1/ots2* (gray square), and *ots1/ots2/phyB-9* (white triangle) seedlings grown in RL is shown. For additional details, see the Fig. 2 legend. (B) The *ots1/ots2* mutant exhibits normal photomorphogenesis in FRL or BL. Hypocotyl elongation inhibition in FRL-grown (Col-0, black triangle; *ots1/ots2*, white triangle) and BL-grown (Col-0, black square; *ots1/ots2*, white square) seedlings is shown. For additional details, see the Fig. 2 legend. (C) Accumulation of the SUMOylated form of phyB is enhanced in transgenic seedlings lacking functional OTS1/OTS2 SUMO proteases. Transgenic *phyB-9* (lane 1) and *ots1/ots2/phyB-9* (lane 2) seedlings expressing *LIP1:PHYB-YFP* were grown under 12 h L/12 h D cycles for 5 d, and samples were harvested on the sixth day at MOD. Accumulation of the SUMOylated phyB-YFP was detected as described in Fig. 1. (D) Recombinant OTS1 deSUMOylates phyB-GFP in vitro. SUMOylated phyB-GFP protein was affinity purified from total protein extracts with anti-GFP antibody and incubated with an identical amount of recombinant OTS1 (His:OTS1) (lane 1) or catalytically inactive OTS1 (His:OTS1<sup>CS</sup>) (lane 2) proteins. After incubation (typically 4 h at room temperature), the beads were reapplied to the column [the flow-through was used to determine 6xHis-OTS1 levels with anti-Histidine tag antibodies (IB: $\alpha$ His)] and washed, and bound proteins were eluted with SDS/PAGE loading buffer and separated by electrophoresis, blotted, and probed with anti-GFP (IB: $\alpha$ GFP) and anti-SUMO1 (IB: $\alpha$ SUMO1) antibodies.

seedlings (Figs. 3 and 4). At present we do not know if this is caused by conjugation of SUMO onto phyB at a second site or whether this is functionally relevant.

SUMOylation of phyB interferes with the conformation-specific interaction of phyB and the negative regulatory factor PIF5 (Fig. 5). We postulate that SUMOylation of phyB at its C-terminal domain directly inhibits interaction of these proteins. This model is supported by recent findings demonstrating that (i) mutations in the C-terminal domain of phyB inhibit binding of PIF3 (53) and that (ii) mutations that fully inactivate signaling by the N-terminal domain of phyB and binding of PIF3 in vitro only partially impair the activity of the full-length photoreceptor (54). Thus, it is plausible to assume that the interaction of PIFs with phyB Pfr is mediated by multiple binding sites and conjugation of the bulky SUMO to the C-terminal domain of phyB substantially weakens its interaction with PIFs. However, it has been recently shown that phosphorylation of phyB<sup>Tyr104</sup> also inhibits binding of PIF3 to phyB Pfr (16), whereas phosphorylation of phyB<sup>Ser86</sup> accelerates thermal reversion (also called D reversion) of phyB Pfr into Pr and thereby prevents interaction of these proteins (15). Thus, it is also possible

that SUMOylation interferes with binding of kinases/phosphatases to phyB and thus indirectly via modifying phosphorylation status of phyB alters the interaction of phyB Pfr with PIFs. Binding of PIFs to phyB Pfr has been shown to be essential for mutual degradation of these proteins (14). Our data illustrate that extended RL treatment moderately decreases the abundance of phyB<sup>Lys996Arg</sup>-YFP compared with phyB-YFP. This observation can be readily explained by compromised interaction of phyB-YFP and PIFs, which is brought about by increased SUMOylation of phyB-YFP compared with the phyB<sup>Lys996Arg</sup>-YFP mutant.

We show that SUMOylation of phyB is a reversible process and propose that the OTS1 and OTS2 SUMO proteases play a significant role in deconjugating SUMO from phyB. This conclusion is supported by data demonstrating that phyB-YFP is hyper-SUMOylated in *ots1/ots2/phyB-9* compared with *phyB-9* (Fig. 6C), OTS1 de-SUMOylates phyB in vitro (Fig. 6D), OTS1 binds to phyB *in planta* (Fig. S8), and the *ots1/ots2* double mutant is hypersensitive to RL (Fig. 6A). However, this mutant exhibits normal BL and FRL-induced physiological responses, indicating that other photoreceptors are not targets of SUMOylation or not deSUMOylated by OTS1/OTS2. SUMOylated phyB is hardly detectable in etiolated or FRL irradiated seedlings, whereas it accumulates to substantially higher levels in RL and WL. These data and the fact that the constitutively active phyB<sup>Tyr276His</sup> mutant is highly SUMOylated both in WL and darkness indicate that SUMOylated phyB preferentially accumulates in its Pfr form. This can be explained either by conformation-specific SUMOylation and/or by de-SUMOylation of phyB. OTS1 binds to phyB Pfr in vivo (Fig. S8), removes SUMO from phyB Pfr in vitro (Fig. 6D), and its abundance is not regulated by light (Fig. S7 C and D). Thus, we conclude that OTS1 is able to de-SUMOylate phyB Pfr as well as phyB Pr, as the abundance of SUMOylated phyB rapidly declines during the D phase in seedlings grown under L/D cycles (Fig. 1B). These data suggest that light-regulated expression and/or modification of yet unknown SUMO proteases or conjugating enzymes or their conformation-specific interaction with phyB Pfr are responsible for the higher level accumulation of SUMOylated phyB Pfr compared with that of phyB Pr.

Independent of the precise mechanism mediating reversible SUMOylation of phyB, we note that SUMOylation has been shown to play a pervasive role in the regulation of stress responses including heat (35), salt (31), and ABA signaling (38). Interestingly, it was shown that OTS1/OTS2 is destabilized by salt stress, and overexpression of OTS1/OTS2 significantly alters salt tolerance of the transgenic plants (31). Because OTS1/OTS2 is involved in mediating reversible SUMOylation of the photoreceptor, we speculate that SUMOylation of phyB might play an important role in the integration of light, salt, and other stress-induced responses.

## Materials and Methods

**Plant Material and Growth Conditions.** Col-0 and *phyB-9* (55) mutant plants were used in this work. Plants expressing PHYB-YFP or N651-GFP in *phyB-9* background have been described (7, 15, 43). The cDNA fragment corresponding to amino acid residues 1–991 of the PHYB protein was inserted as an XbaI-SmaI fragment between the 35S promoter and the YFP-NOS terminator of pPCVB (47). The Lys996Arg and Tyr276His mutations were created by a site-directed mutagenesis kit (Stratagene). Plants were transformed by the Agrobacterium-mediated floral dip method (56). Basta-resistant plants were selected and grown to maturation in the greenhouse. Homozygous T3 plants grown at 22 °C were used in all assays. WL was provided by cool-white fluorescent tubes at a 25  $\mu\text{mol m}^{-2} \text{s}^{-1}$  fluence rate. RL ( $\lambda_{\text{max}}$ , 667 nm) or FRL ( $\lambda_{\text{max}}$ , 730 nm) was provided by LED light sources (Quantum Devices Inc.) at 25 or 10  $\mu\text{mol m}^{-2} \text{s}^{-1}$  fluence rates, respectively. For diurnal conditions, plants were grown in 12 h WL (25  $\mu\text{mol m}^{-2} \text{s}^{-1}$ )/12 h D or 12 h RL (25  $\mu\text{mol m}^{-2} \text{s}^{-1}$ )/12 h D cycles for 7 d before harvesting.

**Transient Assays in *N. benthamiana*.** PIF5 cDNA was cloned in pGWB17 so that expressed proteins had a C-terminal 4xMYC tag, and SUMO1 was cloned in

pEG201 so that expressed proteins had a N-terminal HA tag. Gene constructs were transiently expressed in *N. benthamiana* plants using *Agrobacterium*-mediated transformation (57). The protein samples extracted (57) were mixed with 50  $\mu$ L anti-GFP (Chromotek anti-GFP beads) and incubated on ice for 15 min. The beads were centrifuged down at  $10,000 \times g$  for 1 h and washed three times with 1 mL of cold IP buffer. After the last wash, 50  $\mu$ L of preheated (95 °C) 1 $\times$  SDS loading buffer was used to elute the immuno-complex and analyzed on 8–10% SDS/PAGE using immunoblotting methods with Clontech (clone, JL-8) anti-GFP (Abcam) anti-Myc, and anti-HA antibodies.

**Western Blot Analysis, Hypocotyl Length Measurement.** For details, see *SI Materials and Methods*.

**ACKNOWLEDGMENTS.** Work performed in Edinburgh and Szeged was supported by Biotechnology and Biological Sciences Research Council Grant BB/K006975/1 and Hungarian Scientific Research Fund Grants K-108559 and NN-110636 (to F.N.). A.V. was supported by a Bolyai János Scholarship of the Hungarian Academy of Sciences. Work in the A.S. laboratory in Durham was supported by the Biotechnology and Biological Sciences Research Council and the European Research Council consolidator grant scheme.

- Liu H, Liu B, Zhao C, Pepper M, Lin C (2011) The action mechanisms of plant cryptochromes. *Trends Plant Sci* 16(12):684–691.
- Goyal A, Szarzynska B, Fankhauser C (2013) Phototropism: At the crossroads of light-signaling pathways. *Trends Plant Sci* 18(7):393–401.
- Quail PH (2010) Phytochromes. *Curr Biol* 20(12):R504–R507.
- Rizzini L, et al. (2011) Perception of UV-B by the Arabidopsis UVR8 protein. *Science* 332(6025):103–106.
- Bae G, Choi G (2008) Decoding of light signals by plant phytochromes and their interacting proteins. *Annu Rev Plant Biol* 59:281–311.
- Sharrock RA, Clack T (2002) Patterns of expression and normalized levels of the five Arabidopsis phytochromes. *Plant Physiol* 130(1):442–456.
- Kircher S, et al. (2002) Nucleocytoplasmic partitioning of the plant photoreceptors phytochrome A, B, C, D, and E is regulated differentially by light and exhibits a diurnal rhythm. *Plant Cell* 14(7):1541–1555.
- Chen M, Schwab R, Chory J (2003) Characterization of the requirements for localization of phytochrome B to nuclear bodies. *Proc Natl Acad Sci USA* 100(24):14493–14498.
- Leivar P, Quail PH (2011) PIFs: Pivotal components in a cellular signaling hub. *Trends Plant Sci* 16(1):19–28.
- Huang X, Ouyang X, Deng XW (2014) Beyond repression of photomorphogenesis: Role switching of COP/DET/FUS in light signaling. *Curr Opin Plant Biol* 21:96–103.
- Park E, et al. (2012) Phytochrome B inhibits binding of phytochrome-interacting factors to their target promoters. *Plant J* 72(4):537–546.
- Ma L, et al. (2001) Light control of Arabidopsis development entails coordinated regulation of genome expression and cellular pathways. *Plant Cell* 13(12):2589–2607.
- Leivar P, et al. (2008) The Arabidopsis phytochrome-interacting factor PIF7, together with PIF3 and PIF4, regulates responses to prolonged red light by modulating phyB levels. *Plant Cell* 20(2):337–352.
- Ni W, et al. (2014) A mutually assured destruction mechanism attenuates light signaling in Arabidopsis. *Science* 344(6188):1160–1164.
- Medzihradsky M, et al. (2013) Phosphorylation of phytochrome B inhibits light-induced signaling via accelerated dark reversion in Arabidopsis. *Plant Cell* 25(2):535–544.
- Nito K, Wong CC, Yates JR, 3rd, Chory J (2013) Tyrosine phosphorylation regulates the activity of phytochrome photoreceptors. *Cell Reports* 3(6):1970–1979.
- Johnson ES (2004) Protein modification by SUMO. *Annu Rev Biochem* 73:355–382.
- Saracco SA, Miller MJ, Kurepa J, Vierstra RD (2007) Genetic analysis of SUMOylation in Arabidopsis: Conjugation of SUMO1 and SUMO2 to nuclear proteins is essential. *Plant Physiol* 145(1):119–134.
- Lois LM (2010) Diversity of the SUMOylation machinery in plants. *Biochem Soc Trans* 38(Pt 1):60–64.
- Elrouby N, Coupland G (2010) Proteome-wide screens for small ubiquitin-like modifier (SUMO) substrates identify Arabidopsis proteins implicated in diverse biological processes. *Proc Natl Acad Sci USA* 107(40):17415–17420.
- Song J, Durrin LK, Wilkinson TA, Krutir TG, Chen Y (2004) Identification of a SUMO-binding motif that recognizes SUMO-modified proteins. *Proc Natl Acad Sci USA* 101(40):14373–14378.
- Kerscher O (2007) SUMO junction-what's your function? New insights through SUMO-interacting motifs. *EMBO Rep* 8(6):550–555.
- Hannich JT, et al. (2005) Defining the SUMO-modified proteome by multiple approaches in *Saccharomyces cerevisiae*. *J Biol Chem* 280(6):4102–4110.
- Park HC, et al. (2011) Identification and molecular properties of SUMO-binding proteins in Arabidopsis. *Mol Cells* 32(2):143–151.
- van den Burg HA, Kini RK, Schuurink RC, Takken FL (2010) Arabidopsis small ubiquitin-like modifier paralogs have distinct functions in development and defense. *Plant Cell* 22(6):1998–2016.
- Kurepa J, et al. (2003) The small ubiquitin-like modifier (SUMO) protein modification system in Arabidopsis. Accumulation of SUMO1 and -2 conjugates is increased by stress. *J Biol Chem* 278(9):6862–6872.
- Castro PH, Tavares RM, Bejarano ER, Azevedo H (2012) SUMO, a heavyweight player in plant abiotic stress responses. *Cell Mol Life Sci* 69(19):3269–3283.
- Tomanov K, et al. (2014) Arabidopsis PIAL1 and 2 promote SUMO chain formation as E4-type SUMO ligases and are involved in stress responses and sulfur metabolism. *Plant Cell* 26(11):4547–4560.
- Chosed R, Mukherjee S, Lois LM, Orth K (2006) Evolution of a signalling system that incorporates both redundancy and diversity: Arabidopsis SUMOylation. *Biochem J* 398(3):521–529.
- Colby T, Matthäi A, Boeckelmann A, Stäubli HP (2006) SUMO-conjugating and SUMO-deconjugating enzymes from Arabidopsis. *Plant Physiol* 142(1):318–332.
- Conti L, et al. (2008) Small ubiquitin-like modifier proteases OVERLY TOLERANT TO SALT1 and -2 regulate salt stress responses in Arabidopsis. *Plant Cell* 20(10):2894–2908.
- Ishida T, Yoshimura M, Miura K, Sugimoto K (2012) MMS21/HPY2 and SIZ1, two Arabidopsis SUMO E3 ligases, have distinct functions in development. *PLoS One* 7(10):e46897.
- Murtas G, et al. (2003) A nuclear protease required for flowering-time regulation in Arabidopsis reduces the abundance of SMALL UBIQUITIN-RELATED MODIFIER conjugates. *Plant Cell* 15(10):2308–2319.
- Castañó-Miquel L, et al. (2013) Diversification of SUMO-activating enzyme in Arabidopsis: Implications in SUMO conjugation. *Mol Plant* 6(5):1646–1660.
- Miller MJ, Barrett-Wilt GA, Hua Z, Vierstra RD (2010) Proteomic analyses identify a diverse array of nuclear processes affected by small ubiquitin-like modifier conjugation in Arabidopsis. *Proc Natl Acad Sci USA* 107(38):16512–16517.
- Miura K, et al. (2005) The Arabidopsis SUMO E3 ligase SIZ1 controls phosphate deficiency responses. *Proc Natl Acad Sci USA* 102(21):7760–7765.
- Catala R, et al. (2007) The Arabidopsis E3 SUMO ligase SIZ1 regulates plant growth and drought responses. *Plant Cell* 19(9):2952–2966.
- Miura K, et al. (2009) Sumoylation of ABI5 by the Arabidopsis SUMO E3 ligase SIZ1 negatively regulates abscisic acid signaling. *Proc Natl Acad Sci USA* 106(13):5418–5423.
- Lee J, et al. (2007) Salicylic acid-mediated innate immunity in Arabidopsis is regulated by SIZ1 SUMO E3 ligase. *Plant J* 49(1):79–90.
- Rodríguez MS, Dargemont C, Hay RT (2001) SUMO-1 conjugation in vivo requires both a consensus modification motif and nuclear targeting. *J Biol Chem* 276(16):12654–12659.
- Lamollat F, et al. (2014) Large-scale analysis of lysine SUMOylation by SUMO remnant immunoaffinity profiling. *Nat Commun* 5:5409.
- Ren J, et al. (2009) Systematic study of protein sumoylation: Development of a site-specific predictor of SUMOsp 2.0. *Proteomics* 9(12):3409–3412.
- Palágyi A, et al. (2010) Functional analysis of amino-terminal domains of the photoreceptor phytochrome B. *Plant Physiol* 153(4):1834–1845.
- Kevei E, et al. (2007) Arabidopsis thaliana circadian clock is regulated by the small GTPase LIP1. *Curr Biol* 17(17):1456–1464.
- Khanna R, et al. (2004) A novel molecular recognition motif necessary for targeting photoactivated phytochrome signaling to specific basic helix-loop-helix transcription factors. *Plant Cell* 16(11):3033–3044.
- Kim J, et al. (2003) Functional characterization of phytochrome interacting factor 3 in phytochrome-mediated light signal transduction. *Plant Cell* 15(10):2399–2407.
- Bauer D, et al. (2004) Constitutive photomorphogenesis 1 and multiple photoreceptors control degradation of phytochrome interacting factor 3, a transcription factor required for light signaling in Arabidopsis. *Plant Cell* 16(6):1433–1445.
- Leivar P, Monte E (2014) PIFs: Systems integrators in plant development. *Plant Cell* 26(1):56–78.
- Al-Sady B, Ni W, Kircher S, Schäfer E, Quail PH (2006) Photoactivated phytochrome induces rapid PIF3 phosphorylation prior to proteasome-mediated degradation. *Mol Cell* 23(3):439–446.
- Li L, et al. (2012) Linking photoreceptor excitation to changes in plant architecture. *Genes Dev* 26(8):785–790.
- Matsushita T, Mochizuki N, Nagatani A (2003) Dimers of the N-terminal domain of phytochrome B are functional in the nucleus. *Nature* 424(6948):571–574.
- Krall L, Reed JW (2000) The histidine kinase-related domain participates in phytochrome B function but is dispensable. *Proc Natl Acad Sci USA* 97(14):8169–8174.
- Kircher S, et al. (1999) Light quality-dependent nuclear import of the plant photoreceptors phytochrome A and B. *Plant Cell* 11(8):1445–1456.
- Oka Y, Matsushita T, Mochizuki N, Quail PH, Nagatani A (2008) Mutant screen distinguishes between residues necessary for light-signal perception and signal transfer by phytochrome B. *PLoS Genet* 4(8):e1000158.
- Reed JW, Nagpal P, Poole DS, Furuya M, Chory J (1993) Mutations in the gene for the red/far-red light receptor phytochrome B alter cell elongation and physiological responses throughout Arabidopsis development. *Plant Cell* 5(2):147–157.
- Clough SJ, Bent AF (1998) Floral dip: A simplified method for *Agrobacterium*-mediated transformation of Arabidopsis thaliana. *Plant J* 16(6):735–743.
- Ewan R, et al. (2011) Deubiquitinating enzymes AtUBP12 and AtUBP13 and their tobacco homologue NtUBP12 are negative regulators of plant immunity. *New Phytol* 191(1):92–106.